








# POSITIVELY AWARE 2008 HIV Drug Chart



\*Each drug is recommended for use in combination with other antiretrovirals. Dose may be adjusted when taken in combination. See drug page. Please see package insert for complete information on considerations for indication, efficacy, safety and dose. Not all drugs shown actual size. Drug information current as of November 29, 2007.

Thanks to  
Mariela Diaz-Linares, Pharm.D.


-  Nucleoside and/or Nucleotide Reverse Transcriptase Inhibitor (NRTI, NtRTI)
-  Non-Nucleoside Reverse Transcriptase Inhibitor (NNRTI)
-  Dual-Class Fixed Dosed Combination
-  Protease Inhibitor (PI)
-  Entry Inhibitor
-  Integrase Inhibitor



**COMBIVIR**  
(lamivudine/zidovudine)  
**ADULT SINGLE DOSE\*:**  
One tablet (150 mg lamivudine, 300 mg zidovudine), twice daily  
No food restrictions (with or without food)



**EPZICOM**  
(abacavir/lamivudine)  
**ADULT SINGLE DOSE\*:**  
One tablet (600 mg Ziagen, 300 mg Epivir), once daily  
No food restrictions (with or without food)




**TRUVADA**  
(tenofovir DF/emtricitabine)  
**ADULT SINGLE DOSE\*:**  
One tablet (200 mg Emtriva, 300 mg Viread), once daily  
No food restrictions (with or without food)



**ZERIT**  
(stavudine, d4T)  
**ADULT SINGLE DOSE\*:**  
One 40 mg capsule, twice daily with adjustments for weight (see Drug Guide)  
No food restrictions (with or without food)




**RESCRIPTOR**  
(delavirdine)  
**ADULT SINGLE DOSE\*:**  
Two 200 mg tablets, three times a day or four 100 mg tablets, three times a day  
No food restrictions (with or without food)




**APTIVUS**  
(tipranavir)  
**ADULT SINGLE DOSE\*:**  
Two 250 mg capsules with two 100 mg capsules Norvir, twice daily. Aptivus must be dosed with Norvir.  
Take with food



**KALETRA**  
(lopinavir/ritonavir)  
**ADULT SINGLE DOSE\*:**  
Four tablets (200 mg lopinavir/50 mg ritonavir), once daily if on first-time HIV therapy or two 200/50 mg tablets, twice daily. Half-strength film-coated tablet available: 100 mg lopinavir/25 mg ritonavir.  
No food restrictions (with or without food)




**PREZISTA**  
(darunavir)  
**ADULT SINGLE DOSE\*:**  
Two 300 mg tablets plus 100 mg Norvir, twice daily  
Take with food




**FUZEON**  
(enfuvirtide, T-20)  
**ADULT SINGLE DOSE\*:**  
90 mg, twice daily subcutaneous injections  
No food restrictions (with or without food)




**EMTRIVA**  
(emtricitabine, FTC)  
**ADULT SINGLE DOSE\*:**  
One 200 mg capsule, once daily  
No food restrictions (with or without food)




**RETROVIR**  
(zidovudine, AZT)  
**ADULT SINGLE DOSE\*:**  
One 300 mg tablet, twice daily  
No food restrictions (with or without food)



**VIDEX EC**  
(didanosine, ddi)  
**ADULT SINGLE DOSE\*:**  
One 400 mg capsule, once daily with adjustments for weight (see Drug Guide)  
Take on an empty stomach, at least one hour before or 2 hours after food or drink, except water




**ZIAGEN**  
(abacavir sulfate)  
**ADULT SINGLE DOSE\*:**  
Two 300 mg tablets, once daily (or one 300 mg tablet twice daily)  
No food restrictions (with or without food)




**SUSTIVA**  
(efavirenz)  
**ADULT SINGLE DOSE\*:**  
One 600 mg tablet, once daily or three 200 mg capsules, once daily  
Take on an empty stomach, or may be taken with a light, low-fat snack



**CRIVAN**  
(indinavir)  
**ADULT SINGLE DOSE\*:**  
Two 400 mg capsules, every 8 hours (almost always boosted with Norvir, see Drug Guide)  
Take with no food or a low-fat snack (with food with Norvir); drink at least 1.5 liters of water daily




**LEXIVA**  
(fos-amprenavir calcium)  
**ADULT SINGLE DOSE\*:**  
Two 700 mg tablets plus 100 mg or 200 mg Norvir, once daily or two 700 mg tablets, twice daily (if protease inhibitor experienced, 700 mg with 100 mg Norvir, twice daily)  
No food restrictions (with or without food)




**REYATAZ**  
(atazanavir sulfate)  
**ADULT SINGLE DOSE\*:**  
One 300 mg capsule plus 100 mg Norvir, once daily or two 200 mg capsules, once daily (see Drug Guide)  
Take with food



**SELZENTRY**  
(maraviroc)  
**ADULT SINGLE DOSE\*:**  
150, 300 or 600 mg (available in 150 and 300 mg tablets), twice daily, depends on other medications used (see Drug Guide)  
No food restrictions (with or without food)




**EPIVIR**  
(lamivudine, 3TC)  
**ADULT SINGLE DOSE\*:**  
One 300 mg tablet, once daily; or one 150 mg tablet twice daily  
No food restrictions (with or without food)



**TRIZIVIR**  
(abacavir/lamivudine/zidovudine)  
**ADULT SINGLE DOSE\*:**  
One tablet (300 mg Ziagen, 150 mg Epivir, 300 mg Retrovir), twice daily  
No food restrictions (with or without food)




**VIREAD**  
(tenofovir DF)  
**ADULT SINGLE DOSE\*:**  
One 300 mg tablet, once daily  
No food restrictions (with or without food)



**ATRIPLA**  
(efavirenz/emtricitabine/tenofovir DF)  
**ADULT SINGLE DOSE\*:**  
One tablet (Sustiva [600 mg] and Truvada [200 mg Emtriva, 300 mg Viread]), once daily  
Take on an empty stomach, or may be taken with a light, low-fat snack




**VIRAMUNE**  
(nevirapine)  
**ADULT SINGLE DOSE\*:**  
One 200 mg tablet, daily for the first 14 days, then one 200 mg tablet, twice daily, or two 200 mg tablets, once daily  
No food restrictions (with or without food)




**INVIRASE**  
(saquinavir)  
**ADULT SINGLE DOSE\*:**  
Two 500 mg tablets with one 100 mg capsule Norvir, twice daily. (Or five 200 mg hard-gel capsules and 100 mg Norvir, twice daily.) Saquinavir must be dosed with Norvir.  
Take with food, or within two hours after a meal



**NORVIR**  
(ritonavir)  
**ADULT SINGLE DOSE\*:**  
Used mostly for boosting, once or twice a day with another PI  
Take with food, if possible



**VIRACEPT**  
(nelfinavir)  
**ADULT SINGLE DOSE\*:**  
Two 625 mg tablets, twice daily (or five 250 mg tablets, twice daily)  
Take with a meal



**ISENTRISS**  
(raltegravir, MK-0518)  
**ADULT SINGLE DOSE\*:**  
One 400 mg tablet, twice daily  
No food restrictions (with or without food)